

Alzheimer's Awareness

Everyone is forgetful now and again, and as we age the occurrences may become even more frequent. Yet most people who have mild forgetfulness do not have Alzheimer's disease. But when memory loss starts to interfere with daily activities, it may be time to consult with a physician.

Alzheimer's disease is not a product of the natural aging process. Scientists do not know what causes the disease; however, age is a factor. The disease usually begins after age 60 and the risk goes up with the increase in age. Approximately five million Americans currently suffer from Alzheimer's disease.

According to the Alzheimer's Association (alz.org), there are ten warning signs of Alzheimer's disease:

- ① **Memory loss that disrupts daily life:** forgetting recently learned information or important dates
- ② **Challenges in planning or solving problems:** trouble following familiar recipe or with monthly bills
- ③ **Difficulty completing familiar tasks:** trouble driving to familiar location or managing a budget
- ④ **Confusion with time or place:** losing track of dates, seasons and the passage of time
- ⑤ **Trouble understanding visual images and spatial relationships:** difficulty judging distances and determining color or contrast
- ⑥ **New problems with words or writing:** repeating self, trouble following a conversation
- ⑦ **Misplacing things and inability to retrace steps:** placing things in unusual places and may accuse others of stealing
- ⑧ **Decreased or poor judgment:** problems with money, paying less attention to grooming
- ⑨ **Withdrawal from work or social activities:** trouble completing a favorite hobby and shying away from social engagement
- ⑩ **Changes in mood and personality:** can become confused, suspicious, fearful, anxious and easily upset

Where Quality Caring and Experience Matter • Serving Carbon and surrounding counties for 20 years



23 Main Road • Lehighton, PA 18235

610-377-4443

info@qualitypersonalcare.org
www.QualityPersonalCare.org

The Stages of Alzheimer's

*Developed by Elizabeth Ostuni and Mary Jo Pietro
Lincoln, Nebraska, Chapter, Geriatric Care Managers*

Early "Forgetful" Stages

- Digresses from topic in conversation.
- Tends to repeat self.
- May ramble on and on.
- Relies heavily on clichés.
- Gets along adequately in most social situations.

Middle Stages

- Makes vague, empty, irrelevant conversation.
- Asks fewer questions.
- Is excessively self-oriented.
- Does not initiate conversation.
- Repeats ideas over and over.
- Withdraws from difficult social situations.
- Can still handle some casual social situations.

Late Stages

- Is no longer aware of social interaction or expectancies.
- Withdraws partially or completely from communication.

Although not everyone experiences the same symptoms in the same order or with the same time schedule, we can generally characterize the progress of the disease in six stages, which may last three to 20 years. Note that these represent stages of brain deterioration; they can be caused by diseases other than Alzheimer's.

Forgetfulness

Very mild cognitive decline:

For example, problems such as: subjective complaints about memory deficit such as placement of familiar objects, forgetting names once known well. There is no objective evidence of deficits in social or employment situations. Don't assume that all confusion and memory loss signal Alzheimer's. Reactions to medications can cause reversible delirium and other medical problems may cause dementia. If you are worried, get a medical diagnosis.

Confusion

Early stage: Mild cognitive decline:

For example, problems such as: getting lost when traveling to a familiar location; noticeably lowered performance level at work; trouble finding words and names; little retention from reading; little or no ability to remember names of new people; loss of a valued object; and trouble concentrating.

Late stage: Moderate cognitive decline:

For example, problems such as: decreased knowledge of current and recent events; deficit in memory of personal history; decreased ability to handle travel or finances; and inability to perform complex tasks. Appropriate responsiveness to outside stimulation decreases sharply. Denial of any problem, and withdrawal from challenging situations are common.

Dementia

Early stage: Moderately severe decline:

For example, the person can no longer survive without some assistance. Patients can't remember names of people or places in their lives. They may be disoriented about time and dates. However, they will require no assistance when using the bathroom or eating, but may need help in getting dressed.

Middle stage: Severe cognitive decline:

For example, the person may forget name of spouse and be unaware of events in his or her life. He or she is entirely dependent on others for survival. He or she may have trouble sleeping in a regular pattern.

Late stage: Very severe cognitive decline:

For example, all verbal abilities are lost and he or she needs help eating and using the bathroom. Eventually he or she loses the ability to walk; the brain appears to no longer be able to tell the body what to do.

Based on B. Reisberg, Clinical Presentation, Diagnosis, and Symptomatology of Age-Associated Cognitive Decline and Alzheimer's disease. Alzheimer's Disease: the Standard Reference, New York Free Press, 1983.